

Deltex Medical Bibliography

Pediatrics

Note: The items in [blue](#) are papers where ODM was used or reviewed. Items in black either mention ODM or are documents Deltex Medical are working with to refine the system and develop an understanding of its place in the clinical setting.

Krovetz LJ, McLoughlin TG, Mitchell MB and Schiebler GL. *Hemodynamic Findings in Normal Children*. *Pediat. Res.* 1967. 1: 122-130

Grenadier E, Oliveira C, Allen HD, Sahn DJ, Barron JV, Vales-Cruz LM and Goldberg SJ. *Normal Intracardiac and Great Vessel Doppler Flow Velocities in Infants and Children*. *JACC.* 1984. 4(2): 343-350

Hanseus K, Björkhem G and Lundröm NR. *Cardiac Function in Healthy Infants and Children: Doppler Echocardiographic Evaluation*. *Pediatr Cardiol.* 1994. 15:211-218

Murdoch IA, Marsh MJ, Tibby SM, McLuckie A. *Continuous haemodynamic monitoring in children: use of transoesophageal Doppler*. *Acta Paediatr.* 1995. 84: 761-764

Tibby S, Hatherill M, Marsh MJ and Murdoch IA. *Clinicians' ability to estimate cardiac index in ventilated children and infants*. *Archives of Disease in Childhood.* 1997. 77:516-518

Childs C, Goldring S, Tann W and Hillier VF. *Suprasternal Doppler ultrasound for assessment of stroke distance*. *Arch Dis Child.* 1998. 79:251-255

Tibby SM, Hatherill M, Murdoch IA. *Use of transoesophageal Doppler ultrasonography in ventilated pediatric patients: Derivation of cardiac output*. *Crit Care Med.* 2000. 28:2045-2050

Wodey E, Carre F, Beneux X, Schaffuser A and Ecoffey C. Limits of Corrected Flow Time to Monitor Hemodynamic status in Children. *Journal of Clinical Monitoring and Computing.* 2000. 16: 223-228

Sohn S and Kim HS. Doppler Aortic Flow velocity Measurement in Healthy Children. *J Korean Med.* 2001. 16: 140-4

Tibby SM, Hatherill M, Durward A, Murdoch IA. Are transoesophageal Doppler parameters a reliable guide to paediatric haemodynamic status and fluid management? *Intensive Care Med.* 2001. 27:201-205

Carcillo JA and Fields AI. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Journal de Pediatria.* 2002. 78(6): 449-466

Larousse E, Asehounne K, Datayet B, Albaladejo P, Dubousset AM, Gauthier F and Benhamou D. The hemodynamic effects of pediatric caudal anesthesia assessed by esophageal Doppler. *Anesth Analg.* 2002 May; 94(5): 1165-8

Mohan UR, Britto J, Habibi P, Munter C, Nadel S. Noninvasive Measurement of Cardiac Output in Critically Ill Children. *Journal Pediatric Cardiology.* 2002. 23(1): 58-61

Tibby SM and Murdoch IA. Measurement of cardiac output and tissue perfusion. *Curr Opin Pediatr* 2002, 14:303-309

Chew MS and Poelaert J. Accuracy and repeatability of pediatric cardiac output measurement using Doppler: 20-year review of the literature. *Intensive Care Medicine.* 2003. 29(11): 1889-894

Tibby SM, Murdoch IA. Monitoring cardiac function in intensive care. *Arch Dis Child.* 2003. 88:46-52

King SL and Lim MS. The Use of the Oesophageal Doppler Monitor in the Intensive Care Unit. *Critical Care and Resuscitation.* 2004. 6: 113-122

Raux O, Rochette A, Morau E, Dadure C, Vergnes C and Capdevila X. The effects of spread of block and adrenaline on cardiac output after epidural anesthesia in young children: a randomized, double-blind, prospective study. *Anesth Analg.* 2004. 98(4): 948-55

Hack H. Case report. Use of the Esophageal Doppler Machine to help guide the intraoperative management of two children with pheochromocytoma. *Pediatric Anesthesia*. 2006. 16:867-87

Monsel A, Salvat-Toussaint A, Durand P, Haas V, Baujard C, Rouleau P, El Aouadi S, Benhamou D and Asehnoune K. The Transesophageal Doppler and Hemodynamic Effects of Epidural Anesthesia in Infants Anesthetized with Sevoflurane and Sufentanil. *International Anesthesia Research Society*. 2007. 105(7): 46-50

Rowlands H, Bagshaw O and Duncan H. Does Trans-Oesophageal Doppler Cardiac Output Measurement Change Clinical Management Strategy? 2007 Poster

Knirsch W, Kretschmar O, Tomaske M, Stutz K, Nagdyman N, Balmer C, Schmitz A, Bettex D, Berger F, Bauersfeld U and Weiss M. Cardiac output measurement in children: comparison of the Ultrasound Cardiac Output Monitor with thermodilution cardiac output measurement. *Intensive Care Med*. 2008. 34(6): 1060-4

Mukhtar AM and Obayah G. Esophageal Doppler Monitor: A New Tool in Monitoring Video Assisted Thorascopic Surgery for Ligation of Patent Ductus Arteriosus. *Anesthesia & Analgesia*. 2008. 107(1) 346-347

Schubert S, Schmitz T, Weiss M, Nagdyman N, Huebler M, Alexi-Meskishvili V, Berger F, and Stiller B. Continuous, non-invasive techniques to determine cardiac output in children after cardiac surgery: evaluation of transesophageal Doppler and electric velocimetry. *J Clin Monit Comput*. 2008. 22(4): 299-307

Brierley J et al. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine. *Crit Care Med*. 2009. 37(2) 666-688

Fleck T, Schubert S, Stiller B, Redlin M, Ewert P, Nagdyman N and Berger F, Capability of a new paediatric oesophageal Doppler monitor to detect changes in cardiac output during testing of external pacemakers after cardiac surgery. *J Clin Monit Comput*. 2011. 25(6): 419-25

Absi MA, Lutterman J and Wetzel GT. Noninvasive cardiac output monitoring in the pediatric intensive care unit. *Current Opinion in Cardiology*. 2010. 25: 77-79

Lemson J, Nusmeier A and van de Hoeven JG. Advanced Hemodynamic Monitoring in Critically Ill Children. *Pediatrics*. 2011. 128: 560-571

Lechner E, Hofer A, Leitner-Peneder G, Freynschlag R, Mair R, Weinzettel R, Rehak P and Gombotz H. Levosimendan versus milrinone in neonates and infants after corrective open-heart surgery: a pilot study. *Pediatr Crit Care Med*. 2012 Sep;13(5):542-8.

Brown Z, Gorges M, Cooke E, Malberbe S, Dumont G and Ansermino J. Changes in cardiac index and blood pressure on positioning children prone for scoliosis surgery. *Anaesthesia*. 2013. 68: 742-746

Galante D and Melchionda M. Transesophageal Doppler ultrasound and hemodynamic monitoring using regional anesthesia in pediatrics. *Ultrasounds Anesthesia Journal*. 2013. 1: 1-5

Gan H, Cannesson M, Chandler JR and Ansermino JM. Predicting Fluid Responsiveness in Children: A Systematic Review. *Anesthesia & Analgesia*. 2013. 117(6): 1380- 1392

Dubost C, Bouglé A, Hallynck C, Le Dorze M, Roulleau P, Baujard C and Benhamou D. Comparison of monitoring performance of bioreactance versus esophageal Doppler in pediatric patients. *Indian Journal of Crit Care Med*. 2015 19(1): 3-8

Weber T, Wagner T, Neuman K and Deusch E. Low Predictability of Three Different Noninvasive Methods to determine Fluid responsiveness in Critically Ill Children. *Pediatric Critical Care*. 2015. 16(3): 89-94

Desgranges F P, Desebbe O, Pereira de Souza Neto E, Raphael D, Chassard D. Respiratory variation in aortic blood flow peak velocity to predict fluid responsiveness in mechanically ventilated children: a systematic review and meta-analysis. *Paediatr Anaesth*. 2016 Jan;26(1):37-47

For any enquiries or requests regarding clinical papers on ODM or IOFM, please contact clinical@deltexmedical.com