

## Paediatric Reference List

1. **Tibby et al.** Are transoesophageal Doppler parameters a reliable guide to paediatric haemodynamic status and fluid management? *Intensive Care Med* 2001, 27:201-5.

<http://www.ncbi.nlm.nih.gov/pubmed/11280635>

2. **Tibby et al.** Use of transoesophageal Doppler ultrasonography in ventilated pediatric patients: Derivation of cardiac output. *Crit Care Med* 2000, 28:2045-50.

<http://www.ncbi.nlm.nih.gov/pubmed/10890662>

3. **Lechner et al.** Levosimendan versus milrinone in neonates and infants after corrective open-heart surgery: a pilot study. *Pediatr Crit Care Med* 2012, 13(5):542-8.

<http://www.ncbi.nlm.nih.gov/pubmed/22622649>

4. **Murdoch et al.** Continuous haemodynamic monitoring in children: use of transoesophageal Doppler. *Acta Paediatr* 1995, 84:761-4.

<http://www.ncbi.nlm.nih.gov/pubmed/7549293>

5. **Schubert et al.** Continuous, non-invasive techniques to determine cardiac output in children after cardiac surgery: evaluation of transesophageal Doppler and electric velocimetry. *J Clin Monit Comput* 2008, 22(4):299-307.

<http://www.ncbi.nlm.nih.gov/pubmed/18665449>

6. **Mohan et al.** Noninvasive Measurement of Cardiac Output in Critically Ill Children. *Pediatr Cardiol* 2002, 23(1):58-61.

<http://www.ncbi.nlm.nih.gov/pubmed/11922510>

7. **Lefrant et al.** Training is required to improve the reliability of esophageal Doppler to measure cardiac output in critically ill patients. *Intensive Care Med* 1998;24:347-52.

<http://www.ncbi.nlm.nih.gov/pubmed/9609413>